

## Piśmiennictwo

1. Bellet S. Clinical Disorders of the Heart Beat. 3rd ed. *Lea & Febiger* 1971.
2. Prystowsky EN, Katz AM. Atrial Fibrillation. In: Textbook of Cardiovascular Medicine. *Lippincott-Raven*, Philadelphia 1998: 1661.
3. Knight BP, Michaud GF, Strickberger SA, et al. Electrocardiographic differentiation of atrial flutter from atrial fibrillation by physicians. *J Electrocardiol* 1999; 32: 315-9.
4. Allessie MA, Konings KT, Kirchhof CJ. Mapping of atrial fibrillation. In: Olsson SB, Allessie MA, Campbell RW (eds). Atrial Fibrillation: Mechanisms and Therapeutic Strategies. *Armonk, Futura*, NY 1994: 37-49.
5. Kopecky SL, Gersh BJ, McGoon MD, et al. The natural history of lone atrial fibrillation. A population-based study over three decades. *N Engl J Med* 1987; 317: 669-74.
6. Deleted in proof.
7. Friberg J, Buch P, Scharling H, et al. Rising rates of hospital admissions for atrial fibrillation. *Epidemiology* 2003; 14: 666-72.
8. Le Heuzey JY, Paziaud O, Piot O, et al. Cost of care distribution in atrial fibrillation patients: the COCAF study. *Am Heart J* 2004; 147: 121-6.
9. Go AS, Hylek EM, Phillips KA, et al. Prevalence of diagnosed atrial fibrillation in adults: national implications for rhythm management and stroke prevention: the Anticoagulation and Risk Factors in Atrial Fibrillation (ATRIA) Study. *JAMA* 2001; 285: 2370-5.
10. Furberg CD, Psaty BM, Manolio TA, et al. Prevalence of atrial fibrillation in elderly subjects (the Cardiovascular Health Study). *Am J Cardiol* 1994; 74: 236-41.
11. Friberg J, Scharling H, Gadsbøll N, et al. Sex-specific increase in the prevalence of atrial fibrillation (The Copenhagen City Heart Study). *Am J Cardiol* 2003; 92: 1419-23.
12. Levy S, Maarek M, Coumel P, et al. Characterization of different subsets of atrial fibrillation in general practice in France: the ALFA study. The College of French Cardiologists. *Circulation* 1999; 99: 3028-35.
13. Psaty BM, Manolio TA, Kuller LH, et al. Incidence of and risk factors for atrial fibrillation in older adults. *Circulation* 1997; 96: 2455-61.
14. Crijns HJ, Tjeerdsma G, De Kam PJ, et al. Prognostic value of the presence and development of atrial fibrillation in patients with advanced chronic heart failure. *Eur Heart J* 2000; 21: 1238-45.
15. Madrid AH, Bueno MG, Rebollo JM, et al. Use of irbesartan to maintain sinus rhythm in patients with long-lasting persistent atrial fibrillation: a prospective and randomized study. *Circulation* 2002; 106: 331-6.
16. Wachtell K, Lehto M, Gerdts E, et al. Angiotensin II receptor blockade reduces new-onset atrial fibrillation and subsequent stroke compared to atenolol: the Losartan Intervention For End Point Reduction in Hypertension (LIFE) study. *J Am Coll Cardiol* 2005; 45: 712-9.
17. Atrial Fibrillation Investigators. Risk factors for stroke and efficacy of antithrombotic therapy in atrial fibrillation. Analysis of pooled data from five randomized controlled trials [published erratum appears in Arch Intern Med 1994; 154: 2254]. *Arch Intern Med* 1994; 154: 1449-57.
18. Stewart S, Hart CL, Hole DJ, et al. A population-based study of the long-term risks associated with atrial fibrillation: 20-year follow-up of the Renfrew/Paisley study. *Am J Med* 2002; 113: 359-64.
19. Krahn AD, Manfreda J, Tate RB, et al. The natural history of atrial fibrillation: incidence, risk factors, and prognosis in the Manitoba Follow-Up Study. *Am J Med* 1995; 98: 476-84.
20. Poole-Wilson PA, Swedberg K, Cleland JG, et al. Comparison of carvedilol and metoprolol on clinical outcomes in patients with chronic heart failure in the Carvedilol Or Metoprolol European Trial (COMET): randomised controlled trial. *Lancet* 2003; 362: 7-13.
21. Maggioni AP, Latini R, Carson PE, et al. Valsartan reduces the incidence of atrial fibrillation in patients with heart failure: results from the Valsartan Heart Failure Trial (Val-HeFT). *Am Heart J* 2005; 149: 548-57.
22. Wang TJ, Larson MG, Levy D, et al. Temporal relations of atrial fibrillation and congestive heart failure and their joint influence on mortality: the Framingham Heart Study. *Circulation* 2003; 107: 2920-5.
23. Wolf PA, Abbott RD, Kannel WB. Atrial fibrillation as an independent risk factor for stroke: the Framingham Study. *Stroke* 1991; 22: 983-8.
24. Hart RG, Halperin JL. Atrial fibrillation and thromboembolism: a decade of progress in stroke prevention. *Ann Intern Med* 1999; 131: 688-95.
25. Wolf PA, Dawber TR, Thomas HE Jr, et al. Epidemiologic assessment of chronic atrial fibrillation and risk of stroke: the Framingham study. *Neurology* 1978; 28: 973-7.
26. Bharti S, Lev M. Histology of the normal and diseased atrium. In: Fall RH, Podrid PJ (eds). Atrial Fibrillation: Mechanism and Management. *Raven Press*, NY 1992:15-39.
27. Allessie M, Ausma J, Schotten U. Electrical, contractile and structural remodeling during atrial fibrillation. *Cardiovasc Res* 2002; 54: 230-46.
28. Aime-Sempe C, Folliquet T, Rucker-Martin C, et al. Myocardial cell death in fibrillating and dilated human right atria. *J Am Coll Cardiol* 1999; 34: 1577-86.
29. Polontchouk L, Haefliger JA, Ebelt B, et al. Effects of chronic atrial fibrillation on gap junction distribution in human and rat atria. *J Am Coll Cardiol* 2001; 38: 883-91.
30. Mary-Rabine L, Albert A, Pham TD, et al. The relationship of human atrial cellular electrophysiology to clinical function and ultrastructure. *Circ Res* 1983; 52: 188-99.
31. Bailey GW, Braniff BA, Hancock EW, et al. Relation of left atrial pathology to atrial fibrillation in mitral valvular disease. *Ann Intern Med* 1968; 69: 13-20.
32. Pokharel S, van Geel PP, Sharma UC, et al. Increased myocardial collagen content in transgenic rats overexpressing cardiac angiotensin-converting enzyme is related to enhanced breakdown of N-acetyl-Ser-Asp-Lys-Pro and increased phosphorylation of Smad2/3. *Circulation* 2004; 110: 3129-35.
33. Goette A, Staack T, Rocken C, et al. Increased expression of extracellular signal-regulated kinase and angiotensin-converting enzyme in human atria during atrial fibrillation. *J Am Coll Cardiol* 2000; 35: 1669-77.
34. Kumagai K, Nakashima H, Urata H, et al. Effects of angiotensin II type 1 receptor antagonist on electrical and structural remodeling in atrial fibrillation. *J Am Coll Cardiol* 2003; 41: 2197-204.

35. Verheule S, Wilson E, Everett T, et al. Alterations in atrial electrophysiology and tissue structure in a canine model of chronic atrial dilatation due to mitral regurgitation. *Circulation* 2003; 107: 2615-22.
36. Sanders P, Morton JB, Davidson NC, et al. Electrical remodeling of the atria in congestive heart failure: electrophysiological and electroanatomic mapping in humans. *Circulation* 2003; 108: 1461-8.
37. Jais P, Haissaguerre M, Shah DC, et al. A focal source of atrial fibrillation treated by discrete radiofrequency ablation. *Circulation* 1997; 95: 572-6.
38. Haissaguerre M, Jais P, Shah DC, et al. Spontaneous initiation of atrial fibrillation by ectopic beats originating in the pulmonary veins. *N Engl J Med* 1998; 339: 659-66.
39. Schwartzman D, Bazaz R, Nosbisch J. Common left pulmonary vein: a consistent source of arrhythmogenic atrial ectopy. *J Cardiovasc Electrophysiol* 2004; 15: 560-6.
40. Hsu LF, Jais P, Keane D, et al. Atrial fibrillation originating from persistent left superior vena cava. *Circulation* 2004; 109: 828-32.
41. Chen SA, Tai CT, Yu WC, et al. Right atrial focal atrial fibrillation: electrophysiologic characteristics and radiofrequency catheter ablation. *J Cardiovasc Electrophysiol* 1999; 10: 328-35.
42. Jais P, Hocini M, Macle L, et al. Distinctive electrophysiological properties of pulmonary veins in patients with atrial fibrillation. *Circulation* 2002; 106: 2479-85.
43. Shah D, Haissaguerre M, Jais P, et al. Nonpulmonary vein foci: do they exist? *Pacing Clin Electrophysiol* 2003; 26: 1631-5.
44. Ortiz J, Niwano S, Abe H, et al. Mapping the conversion of atrial flutter to atrial fibrillation and atrial fibrillation to atrial flutter: insights into mechanism. *Circ Res* 1994; 74: 882-94.
45. Konings KT, Kirchof CJ, Smeets JR, et al. High-density mapping of electrically induced atrial fibrillation in humans. *Circulation* 1994; 89: 1665-80.
46. Moe GK, Abildskov JA. Atrial fibrillation as a self sustaining arrhythmia independent of focal discharge. *Am Heart J* 1959; 58: 59-70.
47. Cox JL, Canavan TE, Schuessler RB, et al. The surgical treatment of atrial fibrillation. II. Intraoperative electrophysiologic mapping and description of the electrophysiologic basis of atrial flutter and atrial fibrillation. *J Thorac Cardiovasc Surg* 1991; 101: 406-26.
- 47a. Mandapati R, Skanes A, Chen J, et al. Stable microreentrant sources as a mechanism of atrial fibrillation in the isolated sheep heart. *Circulation* 2000; 101: 194-9.
- 47b. Lazar S, Dixit S, Marchlinski FE, et al. Presence of left-to-right atrial frequency gradient in paroxysmal but not persistent atrial fibrillation in humans. *Circulation* 2004; 110: 3181-6.
- 47c. Sanders P, Berenfeld O, Hocini M, et al. Spectral analysis identifies sites of high-frequency activity maintaining atrial fibrillation in humans. *Circulation* 2005; 112: 789-97.
48. Nakao K, Seto S, Ueyama C, et al. Extended distribution of prolonged and fractionated right atrial electrograms predicts development of chronic atrial fibrillation in patients with idiopathic paroxysmal atrial fibrillation. *J Cardiovasc Electrophysiol* 2002; 13: 996-1002.
49. Akyurek O, Sayin T, Dincer I, et al. Lengthening of intraatrial conduction time in atrial fibrillation and its relation with early recurrence of atrial fibrillation. *Jpn Heart J* 2001; 42: 575-84.
50. Yamada T, Fukunami M, Shimonagata T, et al. Prediction of paroxysmal atrial fibrillation in patients with congestive heart failure: a prospective study. *J Am Coll Cardiol* 2000; 35: 405-13.
51. Ricard P, Levy S, Trigano J, et al. Prospective assessment of the minimum energy needed for external electrical cardioversion of atrial fibrillation. *Am J Cardiol* 1997; 79: 815-6.
52. Wijffels MC, Kirchhof CJ, Dorland R, et al. Atrial fibrillation begets atrial fibrillation. A study in awake chronically instrumented goats. *Circulation* 1995; 92: 1954-68.
53. Nattel S. New ideas about atrial fibrillation 50 years on. *Nature* 2002; 415: 219-26.
54. Anne W, Willems R, Van der Merwe N, et al. Atrial fibrillation after radiofrequency ablation of atrial flutter: preventive effect of angiotensin converting enzyme inhibitors, angiotensin II receptor blockers, and diuretics. *Heart* 2004; 90: 1025-30.
55. Healey JS, Baranchuk A, Crystal E, et al. Prevention of atrial fibrillation with angiotensin-converting enzyme inhibitors and angiotensin receptor blockers: a meta-analysis. *J Am Coll Cardiol* 2005; 45: 1832-9.
56. Pedersen OD, Bagger H, Kober L, et al. Trandolapril reduces the incidence of atrial fibrillation after acute myocardial infarction in patients with left ventricular dysfunction. *Circulation* 1999; 100: 376-80.
57. Prystowsky EN. Atrioventricular node reentry: physiology and radiofrequency ablation. *Pacing Clin Electrophysiol* 1997; 20: 552-71.
58. Page RL, Wharton JM, Prystowsky EN. Effect of continuous vagal enhancement on concealed conduction and refractoriness within the atrioventricular node. *Am J Cardiol* 1996; 77: 260-5.
59. Moe GK, Abildskov JA. Observations on the ventricular dysrhythmia associated with atrial fibrillation in the dog heart. *Circ Res* 1964; 4: 447-60.
60. Van Den Berg MP, Crijns HJ, Haaksma J, et al. Analysis of vagal effects on ventricular rhythm in patients with atrial fibrillation. *Clin Sci (Colch)* 1994; 86: 531-5.
61. Klein GJ, Bashore TM, Sellers TD, et al. Ventricular fibrillation in the Wolff-Parkinson-White syndrome. *N Engl J Med* 1979; 301: 1080-5.
62. Prystowsky EN, Benson DW Jr, Fuster V, et al. Management of patients with atrial fibrillation. A statement for healthcare professionals. From the Subcommittee on Electrocardiography and Electrophysiology, American Heart Association. *Circulation* 1996; 93: 1262-77.
63. Brookes CI, White PA, Staples M, et al. Myocardial contractility is not constant during spontaneous atrial fibrillation in patients. *Circulation* 1998; 98: 1762-8.
64. Sanfilippo AJ, Abascal VM, Sheehan M, et al. Atrial enlargement as a consequence of atrial fibrillation. A prospective echocardiographic study. *Circulation* 1990; 82: 792-7.
65. Gosselink AT, Crijns HJ, Hamer HP, et al. Changes in left and right atrial size after cardioversion of atrial fibrillation: role of mitral valve disease. *J Am Coll Cardiol* 1993; 22: 1666-72.
66. Manning WI, Silverman DI, Katz SE, et al. Impaired left atrial mechanical function after cardioversion: relation to the duration of atrial fibrillation. *J Am Coll Cardiol* 1994; 23: 1535-40.
67. Van Den Berg MP, Tuinenburg AE, van Veldhuisen DJ, et al. Cardioversion of atrial fibrillation in the setting of mild to moderate heart failure. *Int J Cardiol* 1998; 63: 63-70.
68. Packer DL, Bardy GH, Worley SJ, et al. Tachycardia-induced car-

- diomyopathy: a reversible form of left ventricular dysfunction. *Am J Cardiol* 1986; 57: 563-70.
69. Shinbane JS, Wood MA, Jensen DN, et al. Tachycardia-induced cardiomyopathy: a review of animal models and clinical studies. *J Am Coll Cardiol* 1997; 29: 709-15.
70. Halperin JL, Hart RG. Atrial fibrillation and stroke: new ideas, persisting dilemmas. *Stroke* 1988; 19: 937-41.
71. Bogousslavsky J, Van Melle G, Regli F, et al. Pathogenesis of anterior circulation stroke in patients with nonvalvular atrial fibrillation: the Lausanne Stroke Registry. *Neurology* 1990; 40: 1046-50.
72. Miller VT, Rothrock JF, Pearce LA, et al. Ischemic stroke in patients with atrial fibrillation: effect of aspirin according to stroke mechanism. *Stroke Prevention in Atrial Fibrillation Investigators. Neurology* 1993; 43: 32-6.
73. Kanter MC, Tegeler CH, Pearce LA, et al. Carotid stenosis in patients with atrial fibrillation. Prevalence, risk factors, and relationship to stroke in the Stroke Prevention in Atrial Fibrillation Study. *Arch Intern Med* 1994; 154: 1372-7.
74. Hart RG, Halperin JL. Atrial fibrillation and stroke: concepts and controversies. *Stroke* 2001; 32: 803-8.
75. Aschenberg W, Schluter M, Kremer P, et al. Transesophageal twodimensional echocardiography for the detection of left atrial appendage thrombus. *J Am Coll Cardiol* 1986; 7: 163-6.
76. Mugge A, Kuhn H, Nikutta P, et al. Assessment of left atrial appendage function by biplane transesophageal echocardiography in patients with nonrheumatic atrial fibrillation: identification of a subgroup of patients at increased embolic risk. *J Am Coll Cardiol* 1994; 23: 599-607.
77. Manning WJ, Leeman DE, Gotch PJ, et al. Pulsed Doppler evaluation of atrial mechanical function after electrical cardioversion of atrial fibrillation. *J Am Coll Cardiol* 1989; 13: 617-23.
78. Grimm RA, Stewart WJ, Maloney JD, et al. Impact of electrical cardioversion for atrial fibrillation on left atrial appendage function and spontaneous echo contrast: characterization by simultaneous transesophageal echocardiography. *J Am Coll Cardiol* 1993; 22: 1359-66.
79. Chimowitz MI, DeGeorgia MA, Poole RM, et al. Left atrial spontaneous echo contrast is highly associated with previous stroke in patients with atrial fibrillation or mitral stenosis. *Stroke* 1993; 24: 1015-9.
80. Stoddard MF, Dawkins PR, Prince CR, et al. Left atrial appendage thrombus is not uncommon in patients with acute atrial fibrillation and a recent embolic event: a transesophageal echocardiographic study. *J Am Coll Cardiol* 1995; 25: 452-9.
81. Manning WJ, Silverman DI, Waksmonski CA, et al. Prevalence of residual left atrial thrombi among patients with acute thromboembolism and newly recognized atrial fibrillation. *Arch Intern Med* 1995; 155: 2193-8.
82. Khan IA. Atrial stunning: determinants and cellular mechanisms. *Am Heart J* 2003; 145: 787-94.
83. Dunn MI, Marcus JL. Atrial mechanical performance following internal and external cardioversion of atrial fibrillation: its relationship to peripheral embolization and acute cerebrovascular accident. *Chest* 2002; 121: 1-3.
84. Berger M, Schweitzer P. Timing of thromboembolic events after electrical cardioversion of atrial fibrillation or flutter: a retrospective analysis. *Am J Cardiol* 1998; 82: 1545-7, A8.
85. Collins LJ, Silverman DI, Douglas PS, et al. Cardioversion of nonrheumatic atrial fibrillation. Reduced thromboembolic complications with 4 weeks of precardioversion anticoagulation are related to atrial thrombus resolution. *Circulation* 1995; 92: 160-3.
86. Fatkin D, Kelly RP, Feneley MP. Relations between left atrial appendage blood flow velocity, spontaneous echocardiographic contrast and thromboembolic risk in vivo. *J Am Coll Cardiol* 1994; 23: 961-9.
87. Black IW, Chesterman CN, Hopkins AP, et al. Hematologic correlates of left atrial spontaneous echo contrast and thromboembolism in nonvalvular atrial fibrillation. *J Am Coll Cardiol* 1993; 21: 451-7.
88. Yang Y, Gross DG, Li Q, et al. Identification of echocardiographic "smoke" in a bench model with transcranial Doppler ultrasound. *Stroke* 2000; 31: 907-14.
89. Agarwal AK, Venugopalan P. Left atrial spontaneous echo contrast in patients with rheumatic mitral valve stenosis in sinus rhythm: relationship to mitral valve and left atrial measurements. *Int J Cardiol* 2001; 77: 63-8.
90. Black IW. Spontaneous echo contrast: where there's smoke there's fire. *Echocardiography* 2000; 17: 373-82.
91. Sparks PB, Jayaprakash S, Vohra JK, et al. Left atrial "stunning" following radiofrequency catheter ablation of chronic atrial flutter. *J Am Coll Cardiol* 1998; 32: 468-75.
92. Hart RG, Pearce LA, Miller VT, et al. Cardioembolic vs. non-cardioembolic strokes in atrial fibrillation: frequency and effect of antithrombotic agents in the stroke prevention in atrial fibrillation studies. *Cerebrovasc Dis* 2000; 10: 39-43.
93. Blackshear JL, Pearce LA, Hart RG, et al. Aortic plaque in atrial fibrillation: prevalence, predictors, and thromboembolic implications. *Stroke* 1999; 30: 834-40.
94. Fang MC, Singer DE, Chang Y, et al. Gender differences in the risk of ischemic stroke and peripheral embolism in atrial fibrillation: the An-Ticoagulation and Risk factors In Atrial fibrillation (ATRIA) study. *Circulation* 2005; 112: 1687-91.
95. Yoshida M, Nakamura Y, Higashikawa M, Kinoshita M. Predictors of ischemic stroke in non-rheumatic atrial fibrillation. *Int J Cardiol* 1996; 56: 61-70.
96. Hart RG, Pearce LA, McBride R, et al. Factors associated with ischemic stroke during aspirin therapy in atrial fibrillation: analysis of 2012 participants in the SPAF I-III clinical trials. The Stroke Prevention in Atrial Fibrillation (SPAF) Investigators. *Stroke* 1999; 30: 1223-9.
97. Stollberger C, Chnupa P, Kronik G, et al. Transesophageal echocardiography to assess embolic risk in patients with atrial fibrillation. ELAT Study Group. Embolism in Left Atrial Thrombi. *Ann Intern Med* 1998; 128: 630-8.
98. Tsai LM, Lin LJ, Teng JK, et al. Prevalence and clinical significance of left atrial thrombus in nonrheumatic atrial fibrillation. *Int J Cardiol* 1997; 58: 163-9.
99. Rathore SS, Berger AK, Weinfurt KP, et al. Acute myocardial infarction complicated by atrial fibrillation in the elderly: prevalence and outcomes. *Circulation* 2000; 101: 969-74.
100. Goldberg RJ, Yarzebski J, Lessard D, et al. Recent trends in the incidence rates of and death rates from atrial fibrillation complicating initial acute myocardial infarction: a community-wide perspective. *Am Heart J* 2002; 143: 519-27.

101. Prystowsky EN. Tachycardia-induced-tachycardia: a mechanism of initiation of atrial fibrillation. In: DiMarco JP, Prystowsky EN (eds). *Atrial Arrhythmias: State of the Art Armonk*. *Futura*, NY 1995.
102. Brugada R, Tapscott T, Czernuszewicz GZ, et al. Identification of a genetic locus for familial atrial fibrillation. *N Engl J Med* 1997; 336: 905-11.
103. Frost L, Hune LJ, Vestergaard P. Overweight and obesity as risk factors for atrial fibrillation or flutter: the Danish Diet, Cancer, and Health Study. *Am J Med* 2005; 118: 489-95.
104. Wang TJ, Parise H, Levy D, et al. Obesity and the risk of new-onset atrial fibrillation. *JAMA* 2004; 292: 2471-7.
105. Fox CS, Parise H, D'Agostino RB Sr, et al. Parental atrial fibrillation as a risk factor for atrial fibrillation in offspring. *JAMA* 2004; 291: 2851-5.
106. Ellinor PT, Shin JT, Moore RK, et al. Locus for atrial fibrillation maps to chromosome 6q14-16. *Circulation* 2003; 107: 2880-3.
107. Darbar D, Herron KJ, Ballew JD, et al. Familial atrial fibrillation is a genetically heterogeneous disorder. *J Am Coll Cardiol* 2003; 41: 2185-92.
108. Fioranelli M, Piccoli M, Mileto GM, et al. Analysis of heart rate variability five minutes before the onset of paroxysmal atrial fibrillation. *Pacing Clin Electrophysiol* 1999; 22: 743-9.
109. Herweg B, Dalal P, Nagy B, et al. Power spectral analysis of heart period variability of preceding sinus rhythm before initiation of paroxysmal atrial fibrillation. *Am J Cardiol* 1998; 82: 869-74.
110. Coumel P. Neural aspects of paroxysmal atrial fibrillation. In: Falk RH, Podrid PJ (eds). *Atrial Fibrillation: Mechanisms and Management*. Raven Press, NY 1992: 109-25.
111. Maisel WH. Autonomic modulation preceding the onset of atrial fibrillation. *J Am Coll Cardiol* 2003; 42: 1269-70.
112. Fetsch T, Bauer P, Engberding R, et al. Prevention of atrial fibrillation after cardioversion: results of the PAFACT trial. *Eur Heart J* 2004; 25: 1385-94.
113. Israel CW, Gronefeld G, Ehrlich JR, et al. Long-term risk of recurrent atrial fibrillation as documented by an implantable monitoring device: implications for optimal patient care. *J Am Coll Cardiol* 2004; 43: 47-52.
114. Page RL, Wilkinson WE, Clair WK, et al. Asymptomatic arrhythmias in patients with symptomatic paroxysmal atrial fibrillation and paroxysmal supraventricular tachycardia. *Circulation* 1994; 89: 224-7.
115. Kerr CR, Boone J, Connolly SJ, et al. The Canadian Registry of Atrial Fibrillation: a noninterventional follow-up of patients after the first diagnosis of atrial fibrillation. *Am J Cardiol* 1998; 82: 82N-5N.
116. Singh BN, Singh SN, Reda DJ, et al. Amiodarone versus sotalol for atrial fibrillation. *N Engl J Med* 2005; 352: 1861-72.
117. Hamer ME, Blumenthal JA, McCarthy EA, et al. Quality-of-life assessment in patients with paroxysmal atrial fibrillation or paroxysmal supraventricular tachycardia. *Am J Cardiol* 1994; 74: 826-9.
118. Savelieva I, Camm AJ. Clinical relevance of silent atrial fibrillation: prevalence, prognosis, quality of life, and management. *J Interv Card Electrophysiol* 2000; 4: 369-82.
119. Krahn AD, Klein GJ, Kerr CR, et al. How useful is thyroid function testing in patients with recent-onset atrial fibrillation? The Canadian Registry of Atrial Fibrillation Investigators. *Arch Intern Med* 1996; 156: 2221-4.
120. Zabalgoitia M, Halperin JL, Pearce LA, et al. Transesophageal echocardiographic correlates of clinical risk of thromboembolism in nonvalvular atrial fibrillation. *Stroke Prevention in Atrial Fibrillation III Investigators*. *J Am Coll Cardiol* 1998; 31: 1622-6.
121. Healey JS, Crystal E, Lamy A, et al. Left Atrial Appendage Occlusion Study (LAAOS): results of a randomized controlled pilot study of left atrial appendage occlusion during coronary bypass surgery in patients at risk for stroke. *Am Heart J* 2005; 150: 288-93.
122. Sherman DG, Kim SG, Boop BS, et al. Occurrence and characteristics of stroke events in the Atrial Fibrillation Follow-up Investigation of Sinus Rhythm Management (AFFIRM) study. *Arch Intern Med* 2005; 165: 1185-91.
123. Deleted in proof.
124. Van Gelder IC, Hagens VE, Bosker HA, et al. A comparison of rate control and rhythm control in patients with recurrent persistent atrial fibrillation. *N Engl J Med* 2002; 347: 1834-40.
125. Gronefeld GC, Lilienthal J, Kuck KH, et al. Impact of rate versus rhythm control on quality of life in patients with persistent atrial fibrillation. Results from a prospective randomized study. *Eur Heart J* 2003; 24: 1430-6.
126. Carlsson J, Miketic S, Windeler J, et al. Randomized trial of ratecontrol versus rhythm-control in persistent atrial fibrillation: the Strategies of Treatment of Atrial Fibrillation (STAF) study. *J Am Coll Cardiol* 2003; 41: 1690-6.
127. Opolski G, Torbicki A, Kosior DA, et al. Rate control vs. rhythm control in patients with nonvalvular persistent atrial fibrillation: the results of the Polish How to Treat Chronic Atrial Fibrillation (HOT CAFE) Study. *Chest* 2004; 126: 476-86.
128. Wyse DG, Waldo AL, DiMarco JP, et al. A comparison of rate control and rhythm control in patients with atrial fibrillation. *N Engl J Med* 2002; 347: 1825-33.
129. Pelargonio G, Prystowsky EN. Rate versus rhythm control in the management of patients with atrial fibrillation. *Nat Clin Pract Cardiovasc Med* 2005; 2: 514-21.
130. Hohnloser SH, Kuck KH, Lilienthal J. Rhythm or rate control in atrial fibrillation Pharmacological Intervention in Atrial Fibrillation (PIAF): a randomised trial. *Lancet* 2000; 356: 1789-94.
131. Hagens VE, Rancho AV, Van SE, et al. Effect of rate or rhythm control on quality of life in persistent atrial fibrillation. Results from the RateControl Versus Electrical Cardioversion (RACE) Study. *J Am Coll Cardiol* 2004; 43: 241-7.
132. Cooper HA, Bloomfield DA, Bush DE, et al. Relation between achieved heart rate and outcomes in patients with atrial fibrillation (from the Atrial Fibrillation Follow-up Investigation of Rhythm Management [AFFIRM] Study). *Am J Cardiol* 2004; 93: 1247-53.
133. Lonnerholm S, Blomstrom P, Nilsson L, et al. Effects of the maze operation on health-related quality of life in patients with atrial fibrillation. *Circulation* 2000; 101: 2607-11.
134. Oral H, Pappone C, Chugh A, et al. Circumferential pulmonary-vein ablation for chronic atrial fibrillation. *N Engl J Med* 2006; 354: 934-41.
135. Levy S, Ricard P, Lau CP, et al. Multicenter low energy transvenous atrial defibrillation (XAD) trial results in different sub-

- sets of atrial fibrillation. *J Am Coll Cardiol* 1997; 29: 750-5.
136. Lown B, Amarasingham R, Neuman J. New method for terminating cardiac arrhythmias: use of synchronized capacitor discharge. *JAMA* 1962; 182: 548-55.
137. Nerheim P, Birger-Botkin S, Piracha L, et al. Heart failure and sudden death in patients with tachycardia-induced cardiomyopathy and recurrent tachycardia. *Circulation* 2004; 110: 247-52.
138. Petri H, Kafka W, Rudolph W. Discrepant effects of oral and intravenous verapamil on A-V conduction in patients with ventricular preexcitation and atrial fibrillation. *Herz* 1983; 8: 144-52.
139. Wittkampf FH, de Jongste MJ, Lie HI, et al. Effect of right ventricular pacing on ventricular rhythm during atrial fibrillation. *J Am Coll Cardiol* 1988; 11: 539-45.
140. Brignole M, Menozzi C, Gianfranchi L, et al. Assessment of atrioventricular junction ablation and VVIR pacemaker versus pharmacological treatment in patients with heart failure and chronic atrial fibrillation: a randomized, controlled study. *Circulation* 1998; 98: 953-60.
141. Kay GN, Ellenbogen KA, Giudici M, et al. The Ablate and Pace Trial: a prospective study of catheter ablation of the AV conduction system and permanent pacemaker implantation for treatment of atrial fibrillation. APT Investigators. *J Interv Card Electrophysiol* 1998; 2: 121-35.
142. Brignole M, Gianfranchi L, Menozzi C, et al. Assessment of atrioventricular junction ablation and DDDR mode-switching pacemaker versus pharmacological treatment in patients with severely symptomatic paroxysmal atrial fibrillation: a randomized controlled study. *Circulation* 1997; 96: 2617-24.
143. Wood MA, Brown-Mahoney C, Kay GN, et al. Clinical outcomes after ablation and pacing therapy for atrial fibrillation: a meta-analysis. *Circulation* 2000; 101: 1138-44.
144. Williamson BD, Man KC, Daoud E, et al. Radiofrequency catheter modification of atrioventricular conduction to control the ventricular rate during atrial fibrillation [published erratum appears in *N Engl J Med* 1995; 332: 479]. *N Engl J Med* 1994; 331: 910-7.
145. Feld GK, Fleck RP, Fujimura O, et al. Control of rapid ventricular response by radiofrequency catheter modification of the atrioventricular node in patients with medically refractory atrial fibrillation. *Circulation* 1994; 90: 2299-307.
146. Evans GT Jr, Scheinman MM, Bardy G, et al. Predictors of in-hospital mortality after DC catheter ablation of atrioventricular junction. Results of a prospective, international, multi-center study. *Circulation* 1991; 84: 1924-37.
147. Leon AR, Greenberg JM, Kanuru N, et al. Cardiac resynchronization in patients with congestive heart failure and chronic atrial fibrillation: effect of upgrading to biventricular pacing after chronic right ventricular pacing. *J Am Coll Cardiol* 2002; 39: 1258-63.
148. Hart RG, Pearce LA, Rothbart RM, et al. Stroke with intermittent atrial fibrillation: incidence and predictors during aspirin therapy. Stroke Prevention in Atrial Fibrillation Investigators. *J Am Coll Cardiol* 2000; 35: 183-7.
149. Adjusted-dose warfarin versus low-intensity, fixed-dose warfarin plus aspirin for high-risk patients with atrial fibrillation: Stroke Prevention in Atrial Fibrillation III randomised clinical trial. *Lancet* 1996; 348: 633-8.
150. Landefeld CS, Goldman L. Major bleeding in outpatients treated with warfarin: incidence and prediction by factors known at the start of outpatient therapy. *Am J Med* 1989; 87: 144-52.
151. Moulton AW, Singer DE, Haas JS. Risk factors for stroke in patients with nonrheumatic atrial fibrillation: a case-control study. *Am J Med* 1991; 91: 156-61.
152. Gage BF, Waterman AD, Shannon W, et al. Validation of clinical classification schemes for predicting stroke: results from the National Registry of Atrial Fibrillation. *JAMA* 2001; 285: 2864-70.
153. van Walraven WC, Hart RG, Wells GA, et al. A clinical prediction rule to identify patients with atrial fibrillation and a low risk for stroke while taking aspirin. *Arch Intern Med* 2003; 163: 936-43.
154. Patients with nonvalvular atrial fibrillation at low risk of stroke during treatment with aspirin: Stroke Prevention in Atrial Fibrillation III Study. The SPAF III Writing Committee for the Stroke Prevention in Atrial Fibrillation Investigators. *JAMA* 1998; 279: 1273-7.
155. Howitt A, Armstrong D. Implementing evidence based medicine in general practice: audit and qualitative study of antithrombotic treatment for atrial fibrillation. *BMJ* 1999; 318: 1324-7.
156. Biblo LA, Yuan Z, Quan KJ, et al. Risk of stroke in patients with atrial flutter. *Am J Cardiol* 2001; 87: 346-9, A9.
157. Petersen P, Boysen G, Godtfredsen J, et al. Placebo-controlled, randomised trial of warfarin and aspirin for prevention of thromboembolic complications in chronic atrial fibrillation. The Copenhagen AFASAK study. *Lancet* 1989; 1: 175-9.
158. Connolly SJ, Laupacis A, Gent M, et al. Canadian Atrial Fibrillation Anticoagulation (CAFA) Study. *J Am Coll Cardiol* 1991; 18: 349-55.
159. Ezekowitz MD, Bridgers SL, James KE, et al. Warfarin in the prevention of stroke associated with nonrheumatic atrial fibrillation. Veterans Affairs Stroke Prevention in Nonrheumatic Atrial Fibrillation Investigators [published erratum appears in *N Engl J Med* 1993; 328: 148]. *N Engl J Med* 1992; 327: 1406-12.
160. The effect of low-dose warfarin on the risk of stroke in patients with nonrheumatic atrial fibrillation. The Boston Area Anticoagulation Trial for Atrial Fibrillation Investigators. *N Engl J Med* 1990; 323: 1505-11.
161. Stroke Prevention in Atrial Fibrillation Study. Final results. *Circulation* 1991; 84: 527-39.
162. Deleted in proof.
163. Deleted in proof.
164. Secondary prevention in non-rheumatic atrial fibrillation after transient ischaemic attack or minor stroke. EAFT (European Atrial Fibrillation Trial) Study Group. *Lancet* 1993; 342: 1255-62.
165. Hart RG, Benavente O, McBride R, et al. Antithrombotic therapy to prevent stroke in patients with atrial fibrillation: a meta-analysis. *Ann Intern Med* 1999; 131: 492-501.
166. Hylek EM, Singer DE. Risk factors for intracranial hemorrhage in outpatients taking warfarin. *Ann Intern Med* 1994; 120: 897-902.
167. Odén A, Fahlén M, Hart RG. Optimal INR for prevention of stroke and death in atrial fibrillation: a critical appraisal. *Thromb Res* 2006; 117: 493-9.
168. Fihn SD, Callahan CM, Martin DC, et al. The risk for and seve-

- rity of bleeding complications in elderly patients treated with warfarin. The National Consortium of Anticoagulation Clinics. *Ann Intern Med* 1996; 124: 970-9.
169. Fang MC, Chang Y, Hylek EM, et al. Advanced age, anticoagulation intensity, and risk for intracranial hemorrhage among patients taking warfarin for atrial fibrillation. *Ann Intern Med* 2004; 141: 745-52.
170. Hylek EM, Go AS, Chang Y, et al. Effect of intensity of oral anticoagulation on stroke severity and mortality in atrial fibrillation. *N Engl J Med* 2003; 349: 1019-26.
171. Hart RG, Tonarelli SB, Pearce LA. Avoiding central nervous system bleeding during antithrombotic therapy: recent data and ideas. *Stroke* 2005; 36: 1588-93.
172. Albers GW, Diener HC, Frison L, et al. Ximelagatran vs. warfarin for stroke prevention in patients with nonvalvular atrial fibrillation: a randomized trial. *JAMA* 2005; 293: 690-8.
173. Hylek EM, Skates SJ, Sheehan MA, et al. An analysis of the lowest effective intensity of prophylactic anticoagulation for patients with nonrheumatic atrial fibrillation. *N Engl J Med* 1996; 335: 540-6.
174. Gullov AL, Koefoed BG, Petersen P. Bleeding during warfarin and aspirin therapy in patients with atrial fibrillation: the AFASAK 2 study. Atrial Fibrillation Aspirin and Anticoagulation. *Arch Intern Med* 1999; 159: 1322-8.
175. Hellemons BS, Langenberg M, Lodder J, et al. Primary prevention of arterial thromboembolism in non-rheumatic atrial fibrillation in primary care: randomised controlled trial comparing two intensities of coumarin with aspirin. *BMJ* 1999; 319: 958-64.
176. Warfarin versus aspirin for prevention of thromboembolism in atrial fibrillation: Stroke Prevention in Atrial Fibrillation II Study. *Lancet* 1994; 343: 687-91.
177. Diener HC, Cunha L, Forbes C, et al. European Stroke Prevention Study. 2. Dipyridamole and acetylsalicylic acid in the secondary prevention of stroke. *J Neurol Sci* 1996; 143: 1-13.
178. European Stroke Prevention Study. ESPS Group. *Stroke* 1990; 21: 1122-30.
179. Posada IS, Barriales V. Alternate-day dosing of aspirin in atrial fibrillation. LASAF Pilot Study Group. *Am Heart J* 1999; 138: 137-43.
180. Farrell B, Godwin J, Richards S, et al. The United Kingdom transient ischaemic attack (UK-TIA) aspirin trial: final results. *J Neurol Neurosurg Psychiatry* 1991; 54: 1044-54.
181. The efficacy of aspirin in patients with atrial fibrillation. Analysis of pooled data from 3 randomized trials. The Atrial Fibrillation Investigators. *Arch Intern Med* 1997; 157: 1237-40.
182. Deleted in proof.
183. Hart RG, Benavente O, Pearce LA. Increased risk of intracranial hemorrhage when aspirin is combined with warfarin: a meta-analysis and hypothesis. *Cerebrovasc Dis* 1999; 9: 215-7.
184. Stellbrink C, Nixdorff U, Hofmann T, et al. Safety and efficacy of enoxaparin compared with unfractionated heparin and oral anticoagulants for prevention of thromboembolic complications in cardioversion of nonvalvular atrial fibrillation: the Anticoagulation in Cardioversion using Enoxaparin (ACE) trial. *Circulation* 2004; 109: 997-1003.
185. Hirsh J, Warkentin TE, Shaughnessy SG, et al. Heparin and low-molecular-weight heparin: mechanisms of action, pharmacokinetics, dosing, monitoring, efficacy, and safety. *Chest* 2001; 119: 64S-94S.
186. Murray RD, Deitcher SR, Shah A, et al. Potential clinical efficacy and cost benefit of a transesophageal echocardiography-guided low-molecular-weight heparin (exoxaparin) approach to antithrombotic therapy in patients undergoing immediate cardioversion from atrial fibrillation. *J Am Soc Echoangiogr* 2001; 14: 200-8.
187. Stein PD, Alpert JS, Bussey HI, et al. Antithrombotic therapy in patients with mechanical and biological prosthetic heart valves. *Chest* 2001; 119: 220S-7S.
188. Blackshear JL, Johnson WD, Odell JA, et al. Thoracoscopic extracardiac obliteration of the left atrial appendage for stroke risk reduction in atrial fibrillation. *J Am Coll Cardiol* 2003; 42: 1249-52.
189. Ostermayer SH, Reisman M, Kramer PH, et al. Percutaneous left atrial appendage transcatheter occlusion (PLAATO system) to prevent stroke in high-risk patients with non-rheumatic atrial fibrillation: results from the international multi-center feasibility trials. *J Am Coll Cardiol* 2005; 46: 9-14.
190. Halperin JL, Gomberg-Maitland M. Obliteration of the left atrial appendage for prevention of thromboembolism. *J Am Coll Cardiol* 2003; 42: 1259-61.
191. Suttorp MJ, Kingma JH, Jessurun ER, et al. The value of class IC antiarrhythmic drugs for acute conversion of paroxysmal atrial fibrillation or flutter to sinus rhythm. *J Am Coll Cardiol* 1990; 16: 1722-7.
192. Platin EV, Michelson EL, Porterfield JK, et al. Esmolol versus verapamil in the acute treatment of atrial fibrillation or atrial flutter. *Am J Cardiol* 1989; 63: 925-9.
193. Azpitarte J, Alvarez M, Baun O, et al. Value of single oral loading dose of propafenone in converting recent-onset atrial fibrillation. Results of a randomized, double-blind, controlled study. *Eur Heart J* 1997; 18: 1649-54.
194. Kochiadakis GE, Igoumenidis NE, Solomou MC, et al. Efficacy of amiodarone for the termination of persistent atrial fibrillation. *Am J Cardiol* 1999; 83: 58-61.
195. Capucci A, Boriani G, Rubino I, et al. A controlled study on oral propafenone versus digoxin plus quinidine in converting recent onset atrial fibrillation to sinus rhythm. *Int J Cardiol* 1994; 43: 305-13.
196. Vaughan Williams EM. A classification of antiarrhythmic actions reassessed after a decade of new drugs. *J Clin Pharmacol* 1984; 24: 129-47.
197. Falk RH, Pollak A, Singh SN, et al. Intravenous dofetilide, a class III antiarrhythmic agent, for the termination of sustained atrial fibrillation or flutter. Intravenous Dofetilide Investigators. *J Am Coll Cardiol* 1997; 29: 385-90.
198. Norgaard BL, Wachtell K, Christensen PD, et al. Efficacy and safety of intravenously administered dofetilide in acute termination of atrial fibrillation and flutter: a multicenter, randomized, double-blind, placebo-controlled trial. Danish Dofetilide in Atrial Fibrillation and Flutter Study Group. *Am Heart J* 1999; 137: 1062-9.
199. Sedgwick ML, Lip G, Rae AP, et al. Chemical cardioversion of atrial fibrillation with intravenous dofetilide. *Int J Cardiol* 1995; 49: 159-66.

200. Torp-Pedersen C, Moller M, Bloch-Thomsen PE, et al. Dofetilide in patients with congestive heart failure and left ventricular dysfunction. Danish Investigations of Arrhythmia and Mortality on Dofetilide Study Group. *N Engl J Med* 1999; 341: 857-65.
201. Lindeboom JE, Kingma JH, Crijns HJ, et al. Efficacy and safety of intravenous dofetilide for rapid termination of atrial fibrillation and atrial flutter. *Am J Cardiol* 2000; 85: 1031-3.
202. Singh S, Zoble RG, Yellen L, et al. Efficacy and safety of oral dofetilide in converting to and maintaining sinus rhythm in patients with chronic atrial fibrillation or atrial flutter: the symptomatic atrial fibrillation investigative research on dofetilide (SAFIRE-D) study. *Circulation* 2000; 102: 2385-90.
203. Borgeat A, Goy JJ, Maendly R, et al. Flecainide versus quinidine for conversion of atrial fibrillation to sinus rhythm. *Am J Cardiol* 1986; 58: 496-8.
204. Suttorp MJ, Kingma JH, Lie AH, et al. Intravenous flecainide versus verapamil for acute conversion of paroxysmal atrial fibrillation or flutter to sinus rhythm. *Am J Cardiol* 1989; 63: 693-6.
205. Capucci A, Lenzi T, Boriani G, et al. Effectiveness of loading oral flecainide for converting recent-onset atrial fibrillation to sinus rhythm in patients without organic heart disease or with only systemic hypertension. *Am J Cardiol* 1992; 70: 69-72.
206. Donovan KD, Power BM, Hockings BE, et al. Intravenous flecainide versus amiodarone for recent-onset atrial fibrillation. *Am J Cardiol* 1995; 75: 693-7.
207. Botto GL, Bonini W, Broffoni T, et al. Regular ventricular rhythms before conversion of recent onset atrial fibrillation to sinus rhythm. *Pacing Clin Electrophysiol* 1994; 17: 2114-7.
208. Donovan KD, Dobb GJ, Coombs LJ, et al. Reversion of recent-onset atrial fibrillation to sinus rhythm by intravenous flecainide. *Am J Cardiol* 1991; 67: 137-41.
209. Barranco F, Sanchez M, Rodriguez J, et al. Efficacy of flecainide in patients with supraventricular arrhythmias and respiratory insufficiency. *Intensive Care Med* 1994; 20: 42-4.
210. Baldi N, Russo VA, Lenti V, et al. Relation between plasma levels and efficacy of flecainide and propafenone for treatment of atrial fibrillation of recent onset. *New Trends Arrhythmias* 1993; 9: 899-906.
211. Stambler BS, Wood MA, Ellenbogen KA. Antiarrhythmic actions of intravenous ibutilide compared with procainamide during human atrial flutter and fibrillation: electrophysiological determinants of enhanced conversion efficacy. *Circulation* 1997; 96: 4298-306.
212. Guo GB, Ellenbogen KA, Wood MA, et al. Conversion of atrial flutter by ibutilide is associated with increased atrial cycle length variability. *J Am Coll Cardiol* 1996; 27: 1083-9.
213. Volgman AS, Carberry PA, Stambler B, et al. Conversion efficacy and safety of intravenous ibutilide compared with intravenous procainamide in patients with atrial flutter or fibrillation. *J Am Coll Cardiol* 1998; 31: 1414-9.
214. Vos MA, Golitsyn SR, Stangl K, et al. Superiority of ibutilide (a new class III agent) over DL-sotalol in converting atrial flutter and atrial fibrillation. The Ibutilide/Sotalol Comparator Study Group. *Heart* 1998; 79: 568-75.
215. Stambler BS, Wood MA, Ellenbogen KA, et al. Efficacy and safety of repeated intravenous doses of ibutilide for rapid conversion of atrial flutter or fibrillation. Ibutilide Repeat Dose Study Investigators. *Circulation* 1996; 94: 1613-21.
216. Ellenbogen KA, Stambler BS, Wood MA, et al. Efficacy of intravenous ibutilide for rapid termination of atrial fibrillation and atrial flutter: a dose-response study [published erratum appears in *J Am Coll Cardiol* 1996; 28: 1082]. *J Am Coll Cardiol* 1996; 28: 130-6.
217. Bertini G, Conti A, Fradella G, et al. Propafenone versus amiodarone in field treatment of primary atrial tachydysrhythmias. *J Emerg Med* 1990; 8: 15-20.
218. Boriani G, Capucci A, Lenzi T, et al. Propafenone for conversion of recent-onset atrial fibrillation. A controlled comparison between oral loading dose and intravenous administration. *Chest* 1995; 108: 355-8.
219. Boriani G, Biffi M, Capucci A, et al. Oral propafenone to convert recent-onset atrial fibrillation in patients with and without underlying heart disease. A randomized, controlled trial. *Ann Intern Med* 1997; 126: 621-5.
220. Fresco C, Proclemer A, Pavan A, et al. Intravenous propafenone in paroxysmal atrial fibrillation: a randomized, placebo-controlled, doubleblind, multicenter clinical trial. Paroxysmal Atrial Fibrillation Italian Trial (PAFIT)-2 Investigators. *Clin Cardiol* 1996; 19: 409-12.
221. Strobandt R, Stiels B, Hoebrechts R. Propafenone for conversion and prophylaxis of atrial fibrillation. Propafenone Atrial Fibrillation Trial Investigators. *Am J Cardiol* 1997; 79: 418-23.
222. Bellandi F, Cantini F, Pedone T, et al. Effectiveness of intravenous propafenone for conversion of recent-onset atrial fibrillation: a placebo-controlled study. *Clin Cardiol* 1995; 18: 631-4.
223. Bianconi L, Mennuni M, Lukic V, et al. Effects of oral propafenone administration before electrical cardioversion of chronic atrial fibrillation: a placebo-controlled study. *J Am Coll Cardiol* 1996; 28: 700-6.
224. Weiner P, Ganam R, Ganem R, et al. Clinical course of recent-onset atrial fibrillation treated with oral propafenone. *Chest* 1994; 105: 1013-6.
225. Di Benedetto S. Quinidine versus propafenone for conversion of atrial fibrillation to sinus rhythm. *Am J Cardiol* 1997; 80: 518-9.
226. Vita JA, Friedman PL, Cantillon C, et al. Efficacy of intravenous propafenone for the acute management of atrial fibrillation. *Am J Cardiol* 1989; 63: 1275-8.
227. Barroffio R, Tisi G, Guzzini F, et al. A randomised study comparing digoxin and propafenone in the treatment of recent onset atrial fibrillation. *Clin Drug Invest* 1995; 9: 277-83.
228. Galve E, Rius T, Ballester R, et al. Intravenous amiodarone in treatment of recent-onset atrial fibrillation: results of a randomized, controlled study. *J Am Coll Cardiol* 1996; 27: 1079-82.
229. Peuhkurinen K, Niemela M, Ylitalo A, et al. Effectiveness of amiodarone as a single oral dose for recent-onset atrial fibrillation. *Am J Cardiol* 2000; 85: 462-5.
230. Zehender M, Hohnloser S, Muller B, et al. Effects of amiodarone versus quinidine and verapamil in patients with chronic atrial fibrillation: results of a comparative study and a 2-year follow-up. *J Am Coll Cardiol* 1992; 19: 1054-9.
231. Hou ZY, Chang MS, Chen CY, et al. Acute treatment of recent-onset atrial fibrillation and flutter with a tailored dosing regimen of intravenous amiodarone. A randomized, digoxin-controlled study. *Eur Heart J* 1995; 16: 521-8.

232. Opolski G, Stanislawska J, Gorecki A, et al. Amiodarone in restoration and maintenance of sinus rhythm in patients with chronic atrial fibrillation after unsuccessful direct-current cardioversion. *Clin Cardiol* 1997; 20: 337-40.
233. Noc M, Stajer D, Horvat M. Intravenous amiodarone versus verapamil for acute conversion of paroxysmal atrial fibrillation to sinus rhythm. *Am J Cardiol* 1990; 65: 679-80.
234. Tielemans RG, Gosselink AT, Crijns HJ, et al. Efficacy, safety, and determinants of conversion of atrial fibrillation and flutter with oral amiodarone. *Am J Cardiol* 1997; 79: 53-7.
235. Vardas PE, Kochiadakis GE, Igoumenidis NE, et al. Amiodarone as a first-choice drug for restoring sinus rhythm in patients with atrial fibrillation: a randomized, controlled study. *Chest* 2000; 117: 1538-45.
236. Kerin NZ, Faitel K, Naini M. The efficacy of intravenous amiodarone for the conversion of chronic atrial fibrillation. Amiodarone vs. quinidine for conversion of atrial fibrillation. *Arch Intern Med* 1996; 156: 49-53.
237. Hohnloser SH, van de Loo A, Baedeker F. Efficacy and proarrhythmic hazards of pharmacologic cardioversion of atrial fibrillation: prospective comparison of sotalol versus quinidine. *J Am Coll Cardiol* 1995; 26: 852-8.
238. Halinen MO, Huttunen M, Paakkinnen S, et al. Comparison of sotalol with digoxin-quinidine for conversion of acute atrial fibrillation to sinus rhythm (the Sotalol-Digoxin-Quinidine Trial). *Am J Cardiol* 1995; 76: 495-8.
239. Madrid AH, Moro C, Marin-Huerta E, et al. Comparison of flecainide and procainamide in cardioversion of atrial fibrillation. *Eur Heart J* 1993; 14: 1127-31.
240. Deleted in proof.
241. Falk RH, Knowlton AA, Bernard SA, et al. Digoxin for converting recent-onset atrial fibrillation to sinus rhythm. A randomized, double-blinded trial. *Ann Intern Med* 1987; 106: 503-6.
242. Singh S, Saini RK, DiMarco J, et al. Efficacy and safety of sotalol in digitalized patients with chronic atrial fibrillation. The Sotalol Study Group. *Am J Cardiol* 1991; 68: 1227-30.
243. Jordaeens L. Conversion of atrial fibrillation to sinus rhythm and rate control by digoxin in comparison to placebo. *Eur Heart J* 1997; 18: 643-8.
244. Deleted in proof.
245. Intravenous digoxin in acute atrial fibrillation. Results of a randomized, placebo-controlled multicentre trial in 239 patients. The Digitalis in Acute Atrial Fibrillation (DAAF) Trial Group. *Eur Heart J* 1997; 18: 649-54.
246. Sung RJ, Tan HL, Karagounis L, et al. Intravenous sotalol for the termination of supraventricular tachycardia and atrial fibrillation and flutter: a multicenter, randomized, double-blind, placebo-controlled study. Sotalol Multicenter Study Group. *Am Heart J* 1995; 129: 739-48.
247. Nakazawa H, Lythall DA, Noh J, et al. Is there a place for the late cardioversion of atrial fibrillation? A long-term follow-up study of patients with post-thyrotoxic atrial fibrillation. *Eur Heart J* 2000; 21: 327-33.
248. Botto GL, Capucci A, Bonini W, et al. Conversion of recent onset atrial fibrillation to sinus rhythm using a single oral loading dose of propafenone: comparison of two regimens. *Int J Cardiol* 1997; 58: 55-61.
249. Deleted in proof.
250. Pilati G, Lenzi T, Trisolino G, et al. Amiodarone versus quinidine for conversion of recent onset atrial fibrillation to sinus rhythm. *Curr Ther Res* 1991; 49: 140-6.
251. Alboni P, Botto GL, Baldi N, et al. Outpatient treatment of recent-onset atrial fibrillation with the 'pill-in-the-pocket' approach. *N Engl J Med* 2004; 351: 2384-91.
252. Capucci A, Villani GQ, Piepoli MF, et al. The role of oral 1C antiarrhythmic drugs in terminating atrial fibrillation. *Curr Opin Cardiol* 1999; 14: 4-8.
253. Alboni P, Tomasi C, Menozzi C, et al. Efficacy and safety of out-of-hospital self-administered single-dose oral drug treatment in the management of infrequent, well-tolerated paroxysmal supraventricular tachycardia. *J Am Coll Cardiol* 2001; 37: 548-53.
254. Capucci A, Villani GQ, Piepoli MF. Reproducible efficacy of loading oral propafenone in restoring sinus rhythm in patients with paroxysmal atrial fibrillation. *Am J Cardiol* 2003; 92: 1345-7.
255. Khan IA. Single oral loading dose of propafenone for pharmacological cardioversion of recent-onset atrial fibrillation. *J Am Coll Cardiol* 2001; 37: 542-7.
256. Feld GK. Atrial fibrillation. Is there a safe and highly effective pharmacological treatment? *Circulation* 1990; 82: 2248-50.
257. Leitch JW, Klein GJ, Yee R, et al. Prognostic value of electrophysiology testing in asymptomatic patients with Wolff-Parkinson-White pattern [published erratum appears in *Circulation* 1991; 83: 1124]. *Circulation* 1990; 82: 1718-23.
258. Gosselink AT, Crijns HJ, Van Gelder IC, et al. Low-dose amiodarone for maintenance of sinus rhythm after cardioversion of atrial fibrillation or flutter. *JAMA* 1992; 267: 3289-93.
259. Hauser TH, Pinto DS, Josephson ME, et al. Safety and feasibility of a clinical pathway for the outpatient initiation of antiarrhythmic medications in patients with atrial fibrillation or atrial flutter. *Am J Cardiol* 2003; 91: 1437-41.
260. Levy S, Ricard P, Gueunoun M, et al. Low-energy cardioversion of spontaneous atrial fibrillation. Immediate and long-term results. *Circulation* 1997; 96: 253-9.
261. Lesser MF. Safety and efficacy of in-office cardioversion for treatment of supraventricular arrhythmias. *Am J Cardiol* 1990; 66: 1267-8.
262. Joglar JA, Hamdan MH, Ramaswamy K, et al. Initial energy for elective external cardioversion of persistent atrial fibrillation. *Am J Cardiol* 2000; 86: 348-50.
263. Wozakowska-Kaplon B, Janion M, Sielski J, et al. Efficacy of biphasic shock for transthoracic cardioversion of persistent atrial fibrillation: can we predict energy requirements? *Pacing Clin Electrophysiol* 2004; 27: 764-8.
264. Bjerkelund CJ, Orning OM. The efficacy of anticoagulant therapy in preventing embolism related to DC electrical conversion of atrial fibrillation. *Am J Cardiol* 1969; 23: 208-16.
265. Arnold AZ, Mick MJ, Mazurek RP, et al. Role of prophylactic anticoagulation for direct current cardioversion in patients with atrial fibrillation or atrial flutter. *J Am Coll Cardiol* 1992; 19: 851-5.
266. Rabbino MD, Likoff W, Dreifus LS. Complications and limitations of direct current countershock. *JAMA* 1964; 190: 417-20.
267. Lown B, Kleiger R, Williams J. Cardioversion and digitalis drugs: changed threshold to electric shock in digitalized anti-

- mals. *Circ Res* 1965; 17: 519-31.
268. Aberg H, Cullhed I. Direct current countershock complications. *Acta Med Scand* 1968; 183: 415-21.
269. Mancini GB, Goldberger AL. Cardioversion of atrial fibrillation: consideration of embolization, anticoagulation, prophylactic pacemaker, and long-term success. *Am Heart J* 1982; 104: 617-21.
270. Timmermans C, Rodriguez LM, Ayers GM, et al. Effect of electrode length on atrial defibrillation thresholds. *J Cardiovasc Electrophysiol* 1998; 9: 582-7.
271. Tielemans RG, Van Gelder IC, Crijns HJ, et al. Early recurrences of atrial fibrillation after electrical cardioversion: a result of fibrillation-induced electrical remodeling of the atria? *J Am Coll Cardiol* 1998; 31: 167-73.
272. Timmermans C, Rodriguez LM, Smeets JL, et al. Immediate reinitiation of atrial fibrillation following internal atrial defibrillation. *J Cardiovasc Electrophysiol* 1998; 9: 122-8.
273. Rossi M, Lown B. The use of quinidine in cardioversion. *Am J Cardiol* 1967; 19: 234-8.
274. Van Gelder IC, Crijns HJ, van Gilst WH, et al. Prediction of uneventful cardioversion and maintenance of sinus rhythm from direct-current electrical cardioversion of chronic atrial fibrillation and flutter. *Am J Cardiol* 1991; 68: 41-6.
275. Lundstrom T, Ryden L. Chronic atrial fibrillation. Long-term results of direct current conversion. *Acta Med Scand* 1988; 223: 53-9.
276. Niebauer MJ, Brewer JE, Chung MK, et al. Comparison of the rectilinear biphasic waveform with the monophasic damped sine waveform for external cardioversion of atrial fibrillation and flutter. *Am J Cardiol* 2004; 93: 1495-9.
277. Van Gelder IC, Crijns HJ, van Gilst WH, et al. Efficacy and safety of flecainide acetate in the maintenance of sinus rhythm after electrical cardioversion of chronic atrial fibrillation or atrial flutter. *Am J Cardiol* 1989; 64: 1317-21.
278. Crijns HJ, Van Gelder IC, Lie KI. Supraventricular tachycardia mimicking ventricular tachycardia during flecainide treatment. *Am J Cardiol* 1988; 62: 1303-6.
279. Van Gelder IC, Crijns HJ, van Gilst WH, et al. Effects of flecainide on the atrial defibrillation threshold. *Am J Cardiol* 1989; 63: 112-4.
280. Oral H, Souza JJ, Michaud GF, et al. Facilitating transthoracic cardioversion of atrial fibrillation with ibutilide pretreatment. *N Engl J Med* 1999; 340: 1849-54.
281. Li H, Natale A, Tomassoni G, et al. Usefulness of ibutilide in facilitating successful external cardioversion of refractory atrial fibrillation. *Am J Cardiol* 1999; 84: 1096-8, A10.
282. Naccarelli GV, Dell'Orfano JT, Wolbrette DL, et al. Cost-effective management of acute atrial fibrillation: role of rate control, spontaneous conversion, medical and direct current cardioversion, transesophageal echocardiography, and antiembolic therapy. *Am J Cardiol* 2000; 85: 36D-45D.
283. Manning WJ, Silverman DI, Gordon SP, et al. Cardioversion from atrial fibrillation without prolonged anticoagulation with use of transesophageal echocardiography to exclude the presence of atrial thrombi. *N Engl J Med* 1993; 328: 750-5.
284. Black IW, Fatkin D, Sagar KB, et al. Exclusion of atrial thrombus by transesophageal echocardiography does not preclude embolism after cardioversion of atrial fibrillation. A multicenter study. *Circulation* 1994; 89: 2509-13.
285. Moreyra E, Finkelhor RS, Cebul RD. Limitations of transesophageal echocardiography in the risk assessment of patients before nonanticoagulated cardioversion from atrial fibrillation and flutter: an analysis of pooled trials. *Am Heart J* 1995; 129: 71-5.
286. Fatkin D, Kuchar DL, Thorburn CW, et al. Transesophageal echocardiography before and during direct current cardioversion of atrial fibrillation: evidence for "atrial stunning" as a mechanism of thromboembolic complications. *J Am Coll Cardiol* 1994; 23: 307-16.
287. Antonielli E, Pizzati A, Bassignana A, et al. Transesophageal echocardiographic evidence of more pronounced left atrial stunning after chemical (propafenone) rather than electrical attempts at cardioversion from atrial fibrillation. *Am J Cardiol* 1999; 84: 1092-10.
288. Falcone RA, Morady F, Armstrong WF. Transesophageal echocardiographic evaluation of left atrial appendage function and spontaneous contrast formation after chemical or electrical cardioversion of atrial fibrillation. *Am J Cardiol* 1996; 78:435-9.
289. Bellotti P, Spirito P, Lupi G, et al. Left atrial appendage function assessed by transesophageal echocardiography before and on the day after elective cardioversion for nonvalvular atrial fibrillation. *Am J Cardiol* 1998; 81: 1199-202.
290. Harjai K, Mobarek S, Abi-Samra F, et al. Mechanical dysfunction of the left atrium and the left atrial appendage following cardioversion of atrial fibrillation and its relation to total electrical energy used for cardioversion. *Am J Cardiol* 1998; 81: 1125-9.
291. Mitusch R, Garbe M, Schmucker G, et al. Relation of left atrial appendage function to the duration and reversibility of nonvalvular atrial fibrillation. *Am J Cardiol* 1995; 75: 944-7.
292. Manning WJ, Silverman DI, Katz SE, et al. Temporal dependence of the return of atrial mechanical function on the mode of cardioversion of atrial fibrillation to sinus rhythm. *Am J Cardiol* 1995; 75: 624-6.
293. Grimm RA, Leung DY, Black IW, et al. Left atrial appendage "stunning" after spontaneous conversion of atrial fibrillation demonstrated by transesophageal Doppler echocardiography. *Am Heart J* 1995; 130: 174-6.
294. Klein AL, Grimm RA, Murray RD, et al. Use of transesophageal echocardiography to guide cardioversion in patients with atrial fibrillation. *N Engl J Med* 2001; 344: 1411-20.
295. Mehta D, Baruch L. Thromboembolism following cardioversion of "common" atrial flutter. Risk factors and limitations of transesophageal echocardiography. *Chest* 1996; 110: 1001-3.
296. Irani WN, Grayburn PA, Afridi I. Prevalence of thrombus, spontaneous echo contrast, and atrial stunning in patients undergoing cardioversion of atrial flutter. A prospective study using transesophageal echocardiography. *Circulation* 1997; 95: 962-6.
297. Lazzeroni E, Picano E, Morozzi L, et al. Dipyridamole-induced ischemia as a prognostic marker of future adverse cardiac events in adult patients with hypertrophic cardiomyopathy. Echo Persantine Italian Cooperative (EPIC) Study Group, Sub-project Hypertrophic Cardiomyopathy. *Circulation* 1997; 96: 4268-72.
298. Deleted in proof.
299. Kerr CR, Humphries KH, Talajic M, et al. Progression to chro-

- nic atrial fibrillation after the initial diagnosis of paroxysmal atrial fibrillation: results from the Canadian Registry of Atrial Fibrillation. *Am Heart J* 2005; 149: 489-96.
300. Van Gelder IC, Crijns HJ, Tielemans RG, et al. Chronic atrial fibrillation. Success of serial cardioversion therapy and safety of oral anticoagulation. *Arch Intern Med* 1996; 156: 2585-92.
301. Van Gelder IC, Tuinenburg AE, Schoonderwoerd BS, et al. Pharmacologic versus direct-current electrical cardioversion of atrial flutter and fibrillation. *Am J Cardiol* 1999; 84: 147R-51R.
302. Suttorp MJ, Kingma JH, Koomen EM, et al. Recurrence of paroxysmal atrial fibrillation or flutter after successful cardioversion in patients with normal left ventricular function. *Am J Cardiol* 1993; 71: 710-3.
303. Prystowsky EN. Management of atrial fibrillation: therapeutic options and clinical decisions. *Am J Cardiol* 2000; 85: 3-11.
304. Singh SN, Fletcher RD, Fisher SG, et al. Amiodarone in patients with congestive heart failure and asymptomatic ventricular arrhythmia. Survival Trial of Antiarrhythmic Therapy in Congestive Heart Failure. *N Engl J Med* 1995; 333: 77-82.
305. Ehrlich JR, Nattel S, Hohnloser SH. Atrial fibrillation and congestive heart failure: specific considerations at the intersection of two common and important cardiac disease sets. *J Cardiovasc Electrophysiol* 2002; 13: 399-405.
306. Maisel WH, Stevenson LW. Atrial fibrillation in heart failure: epidemiology, pathophysiology, and rationale for therapy. *Am J Cardiol* 2003; 91: 2D-8D.
307. Naccarelli GV, Hynes BJ, Wolbrette DL, et al. Atrial fibrillation in heart failure: prognostic significance and management. *J Cardiovasc Electrophysiol* 2003; 14: S281-6.
308. Meng F, Yoshikawa T, Baba A, et al. Beta-blockers are effective in congestive heart failure patients with atrial fibrillation. *J Card Fail* 2003; 9: 398-403.
309. Steeds RP, Birchall AS, Smith M, et al. An open label, randomised, crossover study comparing sotalol and atenolol in the treatment of symptomatic paroxysmal atrial fibrillation. *Heart* 1999; 82: 170-5.
310. Kuhlkamp V, Schirdewan A, Stangl K, et al. Use of metoprolol CR/XL to maintain sinus rhythm after conversion from persistent atrial fibrillation: a randomized, double-blind, placebo-controlled study. *J Am Coll Cardiol* 2000; 36: 139-46.
311. Essebag V, Hadjis T, Platt RW, et al. Amiodarone and the risk of bradyarrhythmia requiring permanent pacemaker in elderly patients with atrial fibrillation and prior myocardial infarction. *J Am Coll Cardiol* 2003; 41: 249-54.
312. Julian DG, Camm AJ, Frangin G, et al. Randomised trial of effect of amiodarone on mortality in patients with left-ventricular dysfunction after recent myocardial infarction: EMIAT. European Myocardial Infarct Amiodarone Trial Investigators [published errata appear in *Lancet* 1997; 349: 1180 and 1997; 349: 1776]. *Lancet* 1997; 349: 667-74.
313. Cairns JA, Connolly SJ, Roberts R, et al. Randomised trial of outcome after myocardial infarction in patients with frequent or repetitive ventricular premature depolarisations: CAMIAT. Canadian Amiodarone Myocardial Infarction Arrhythmia Trial Investigators [published erratum appears in *Lancet* 1997; 349: 1776]. *Lancet* 1997; 349: 675-82.
314. Kober L, Bloch Thomsen PE, Moller M, et al. Effect of dofetilide in patients with recent myocardial infarction and left-ventricular dysfunction: a randomised trial. *Lancet* 2000; 356: 2052-8.
315. Peters NS, Schilling RJ, Kanagaratnam P, et al. Atrial fibrillation: strategies to control, combat, and cure. *Lancet* 2002; 359: 593-603.
316. Tsang TS, Petty GW, Barnes ME, et al. The prevalence of atrial fibrillation in incident stroke cases and matched population controls in Rochester, Minnesota: changes over three decades. *J Am Coll Cardiol* 2003; 42: 93-100.
317. Jackman WM, Friday KJ, Anderson JL, et al. The long QT syndromes: a critical review, new clinical observations and a unifying hypothesis. *Prog Cardiovasc Dis* 1988; 31: 115-72.
318. Van Noord T, Tielemans RG, Bosker HA, et al. Beta-blockers prevent subacute recurrences of persistent atrial fibrillation only in patients with hypertension. *Europace* 2004; 6: 343-50.
319. Klingbeil AU, Schneider M, Martus P, et al. A meta-analysis of the effects of treatment on left ventricular mass in essential hypertension. *Am J Med* 2003; 115: 41-6.
320. L'Allier PL, Ducharme A, Keller PF, et al. Angiotensin-converting enzyme inhibition in hypertensive patients is associated with a reduction in the occurrence of atrial fibrillation. *J Am Coll Cardiol* 2004; 44: 159-64.
321. Yusuf S, Sleight P, Pogue J, et al. Effects of an angiotensin-converting-enzyme inhibitor, ramipril, on cardiovascular events in high-risk patients. The Heart Outcomes Prevention Evaluation Study Investigators. *N Engl J Med* 2000; 342: 145-53.
322. Bosch J, Yusuf S, Pogue J, et al. Use of ramipril in preventing stroke: double blind randomised trial. *BMJ* 2002; 324: 699-702.
323. Chapman N, Huxley R, Anderson C, et al. Effects of a perindopril-based blood pressure-lowering regimen on the risk of recurrent stroke according to stroke subtype and medical history: the PROGRESS Trial. *Stroke* 2004; 35: 116-21.
324. Dahlöf B, Devereux RB, Kjeldsen SE, et al. Cardiovascular morbidity and mortality in the Losartan Intervention For Endpoint reduction in hypertension study (LIFE): a randomised trial against atenolol. *Lancet* 2002; 359: 995-1003.
325. Dahlöf B, Zanchetti A, Diez J, et al. Effects of losartan and atenolol on left ventricular mass and neurohormonal profile in patients with essential hypertension and left ventricular hypertrophy. *J Hypertens* 2002; 20: 1855-64.
326. Arima H, Hart RG, Colman S, et al. Perindopril-based blood pressure-lowering reduces major vascular events in patients with atrial fibrillation and prior stroke or transient ischemic attack. *Stroke* 2005; 36: 2164-9.
327. Wachtell K, Hornestam B, Lehto M, et al. Cardiovascular morbidity and mortality in hypertensive patients with a history of atrial fibrillation: the Losartan Intervention For End Point Reduction in Hypertension (LIFE) study. *J Am Coll Cardiol* 2005; 45: 705-11.
328. Cox JL, Schuessler RB, Lappas DG, et al. An 8 1/2-year clinical experience with surgery for atrial fibrillation. *Ann Surg* 1996; 224: 267-73.
329. Cox JL. Cardiac surgery for arrhythmias. *J Cardiovasc Electrophysiol* 2004; 15: 250-62.
330. Cox JL, Boineau JP, Schuessler RB, et al. Modification of the maze procedure for atrial flutter and atrial fibrillation. I. Rationale and surgical results. *J Thorac Cardiovasc Surg* 1995; 110: 473-84.

331. Damiano RJ Jr, Gaynor SL, Bailey M, et al. The long-term outcome of patients with coronary disease and atrial fibrillation undergoing the Cox maze procedure. *J Thorac Cardiovasc Surg* 2003; 126: 2016-21.
332. Gillinov AM, McCarthy PM. Advances in the surgical treatment of atrial fibrillation. *Cardiol Clin* 2004; 22: 147-57.
333. Packer DL, Asirvatham S, Munger TM. Progress in nonpharmacologic therapy of atrial fibrillation. *J Cardiovasc Electrophysiol* 2003; 14: S296-S309.
334. Chen SA, Hsieh MH, Tai CT, et al. Initiation of atrial fibrillation by ectopic beats originating from the pulmonary veins: electrophysiological characteristics, pharmacological responses, and effects of radiofrequency ablation. *Circulation* 1999; 100: 1879-86.
335. Lin WS, Tai CT, Hsieh MH, et al. Catheter ablation of paroxysmal atrial fibrillation initiated by non-pulmonary vein ectopy. *Circulation* 2003; 107: 3176-83.
336. Hocini M, Sanders P, Jais P, et al. Techniques for curative treatment of atrial fibrillation. *J Cardiovasc Electrophysiol* 2004; 15: 1467-71.
337. Haissaguerre M, Shah DC, Jais P, et al. Electrophysiological breakthroughs from the left atrium to the pulmonary veins. *Circulation* 2000; 102: 2463-5.
338. Verma A, Marrouche NF, Natale A. Pulmonary vein antrum isolation: intracardiac echocardiography-guided technique. *J Cardiovasc Electrophysiol* 2004; 15: 1335-40.
339. Wazni OM, Marrouche NF, Martin DO, et al. Radiofrequency ablation vs. antiarrhythmic drugs as first-line treatment of symptomatic atrial fibrillation: a randomized trial. *JAMA* 2005; 293: 2634-40.
340. Pappone C, Rosanio S, Oreto G, et al. Circumferential radiofrequency ablation of pulmonary vein ostia: a new anatomic approach for curing atrial fibrillation. *Circulation* 2000; 102: 2619-28.
341. Pappone C, Santinelli V. The who, what, why, and how-to guide for circumferential pulmonary vein ablation. *J Cardiovasc Electrophysiol* 2004; 15: 1226-30.
342. Oral H, Scharf C, Chugh A, et al. Catheter ablation for paroxysmal atrial fibrillation: segmental pulmonary vein ostial ablation versus left atrial ablation. *Circulation* 2003; 108: 2355-60.
343. Cappato R, Calkins H, Chen SA, et al. Worldwide survey on the methods, efficacy, and safety of catheter ablation for human atrial fibrillation. *Circulation* 2005; 111: 1100-5.
344. Nademanee K, McKenzie J, Kosar E, et al. A new approach for catheter ablation of atrial fibrillation: mapping of the electrophysiologic substrate. *J Am Coll Cardiol* 2004; 43: 2044-53.
345. Hsu LF, Jais P, Sanders P, et al. Catheter ablation for atrial fibrillation in congestive heart failure. *N Engl J Med* 2004; 351: 2373-83.
346. Pappone C, Rosanio S, Augello G, et al. Mortality, morbidity, and quality of life after circumferential pulmonary vein ablation for atrial fibrillation: outcomes from a controlled nonrandomized longterm study. *J Am Coll Cardiol* 2003; 42: 185-97.
347. Marshall HJ, Harris ZI, Griffith MJ, et al. Prospective randomized study of ablation and pacing versus medical therapy for paroxysmal atrial fibrillation: effects of pacing mode and mode-switch algorithm. *Circulation* 1999; 99: 1587-92.
348. Natale A, Zimerman L, Tomassoni G, et al. AV node ablation and pacemaker implantation after withdrawal of effective rate-control medications for chronic atrial fibrillation: effect on quality of life and exercise performance. *Pacing Clin Electrophysiol* 1999; 22: 1634-9.
349. Marshall HJ, Harris ZI, Griffith MJ, et al. Atrioventricular nodal ablation and implantation of mode switching dual chamber pacemakers: effective treatment for drug refractory paroxysmal atrial fibrillation. *Heart* 1998; 79: 543-7.
350. Hindricks G, Piorkowski C, Tanner H, et al. Perception of atrial fibrillation before and after radiofrequency catheter ablation: relevance of asymptomatic arrhythmia recurrence. *Circulation* 2005; 112: 307-13.
351. Senatore G, Stabile G, Bertaglia E, et al. Role of transtelephonic electrocardiographic monitoring in detecting short-term arrhythmia recurrences after radiofrequency ablation in patients with atrial fibrillation. *J Am Coll Cardiol* 2005; 45: 873-6.
352. Karch MR, Zrenner B, Deisenhofer I, et al. Freedom from atrial tachyarrhythmias after catheter ablation of atrial fibrillation: a randomized comparison between 2 current ablation strategies. *Circulation* 2005; 111: 2875-80.
353. Haissaguerre M, Jais P, Shah DC, et al. Electrophysiological end point for catheter ablation of atrial fibrillation initiated from multiple pulmonary venous foci. *Circulation* 2000; 101: 1409-17.
354. Ren JF, Marchlinski FE, Callans DJ, et al. Increased intensity of anticoagulation may reduce risk of thrombus during atrial fibrillation ablation procedures in patients with spontaneous echo contrast. *J Cardiovasc Electrophysiol* 2005; 16: 474-7.
355. Pappone C, Oral H, Santinelli V, et al. Atrio-esophageal fistula as a complication of percutaneous transcatheter ablation of atrial fibrillation. *Circulation* 2004; 109: 2724-6.
356. Scanavacca MI, D'avila A, Parga J, et al. Left atrial-esophageal fistula following radiofrequency catheter ablation of atrial fibrillation. *J Cardiovasc Electrophysiol* 2004; 15: 960-2.
357. Mesas CE, Pappone C, Lang CC, et al. Left atrial tachycardia after circumferential pulmonary vein ablation for atrial fibrillation: electroanatomic characterization and treatment. *J Am Coll Cardiol* 2004; 44: 1071-9.
358. Pappone C, Manguso F, Vicedomini G, et al. Prevention of iatrogenic atrial tachycardia after ablation of atrial fibrillation: a prospective randomized study comparing circumferential pulmonary vein ablation with a modified approach. *Circulation* 2004; 110: 3036-42.
359. Andersen HR, Nielsen JC, Thomsen PE, et al. Long-term follow-up of patients from a randomised trial of atrial versus ventricular pacing for sick-sinus syndrome. *Lancet* 1997; 350: 1210-6.
360. Connolly SJ, Kerr CR, Gent M, et al. Effects of physiologic pacing versus ventricular pacing on the risk of stroke and death due to cardiovascular causes. Canadian Trial of Physiologic Pacing Investigators. *N Engl J Med* 2000; 342: 1385-91.
361. Lamas GA, Orav EJ, Stambler BS, et al. Quality of life and clinical outcomes in elderly patients treated with ventricular pacing as compared with dual-chamber pacing. Pacemaker Selection in the Elderly Investigators. *N Engl J Med* 1998; 338: 1097-104.

362. Lamas GA, Lee KL, Sweeney MO, et al. Ventricular pacing or dualchamber pacing for sinus-node dysfunction. *N Engl J Med* 2002; 346: 1854-62.
363. Knight BP, Gersh BJ, Carlson MD, et al. Role of permanent pacing to prevent atrial fibrillation: science advisory from the American Heart Association Council on Clinical Cardiology (Subcommittee on Electrocardiography and Arrhythmias) and the Quality of Care and Outcomes Research Interdisciplinary Working Group, in collaboration with the Heart Rhythm Society. *Circulation* 2005; 111: 240-3.
364. Deleted in proof.
365. Blanc JJ, De Roy L, Mansouriati J, et al. Atrial pacing for prevention of atrial fibrillation: assessment of simultaneously implemented algorithms. *Europace* 2004; 6: 371-9.
366. Friedman PA, Ip JH, Jazayeri M, et al. The impact of atrial prevention and termination therapies on atrial tachyarrhythmia burden in patients receiving a dual-chamber defibrillator for ventricular arrhythmias. *J Interv Card Electrophysiol* 2004; 10: 103-10.
367. Alsheikh-Ali AA, Wang PJ, Rand W, et al. Enalapril treatment and hospitalization with atrial tachyarrhythmias in patients with left ventricular dysfunction. *Am Heart J* 2004; 147: 1061-5.
368. Olsson LG, Swedberg K, Ducharme A, et al. on behalf of the CHARM Investigators. Atrial fibrillation and risk of clinical events in chronic heart failure with and without left ventricular systolic dysfunction. Results from the Candesartan in Heart failure-Assessment of Reduction in Mortality and morbidity (CHARM) program. *J Am Coll Cardiol* 2006; 47: 1997-2004.
369. Young-Xu Y, Jabbour S, Goldberg R, et al. Usefulness of statin drugs in protecting against atrial fibrillation in patients with coronary artery disease. *Am J Cardiol* 2003; 92: 1379-83.
370. Siu CW, Lau CP, Tse HF. Prevention of atrial fibrillation recurrence by statin therapy in patients with lone atrial fibrillation after successful cardioversion. *Am J Cardiol* 2003; 92: 1343-5.
371. Mozaffarian D, Psaty BM, Rimm EB, et al. Fish intake and risk of incident atrial fibrillation. *Circulation* 2004; 110: 368-73.